References

- **1.Wilson JJ. & Furukawa M.(2014):** Evaluation of the patient with hip pain. Am Fam Physician. 89(1):27-34.
- 2.Kamal D., Trăistaru R. & Alexandru D.(2013):

 Epidemiologic Study of Avascular Necrosis of the
 Femoral Head. Current Health Sciences Journal
 Vol. 39, No. 3: 169-74.
- **3.Kaushik AP., Das A., & Cui Q. (2012):** Osteonecrosis of the femoral head: An update in year 2012. World Journal of Orthopedics, 3(5), 49–57.
- 4.Weishaupt D., Exner G., Hilfiker P., et al.(2000):

 Dynamic MR Imaging of the Hip in Legg-CalvéPerthes Disease: Comparison with Arthrography.
 In AJR. 174:1635-37.
- **5.Beltran J., Patnana M., Beltran L., et al.(2002):** MRI of the hip. In Applied Radiology. 33-39.
- **6.Boutry N., Paul C., Leroy X., et al.(2002):** Rapidly destructive osteoarthritis of the hip: MR imaging findings. AJR Am J Roentgenol. 179:657-663.
- **7.McWalter P. & Hassan A. (2009):** Transient osteoporosis of the hip. Annals of Saudi Medicine, 29(2), 146–148.
- 8.Bos CF., Bloem JL., Obermann WR., et al (1988):

 Magnetic resonance imaging in congenital

dislocation of the hip. J Bone Joint Surg 70:174–178

- 9.Druschel C., Placzek R., Selka L., et al.(2013): MRI evaluation of hip containment and congruency after closed reduction in congenital hip dislocation. In Hip Int. 23(6): 552 559.
- **10. Lee MC. & Eberson CP.(2006):** Growth and development of the child's hip. Orthop Clin North Am. ;37(2):119-32.
- 11. Karl JJ. & Davies AM.(2006): Congenital and Developmental Abnormalities, In Davies A.M., Johnson K., Richard W. et al. First edition. Imaging of the Hip & Bony Pelvis; Techniques and Applications. Springer. pp. 93-104.
- 12. Ludwig O., Pierre B. &Herman J.(2003): The Hip In a System of Orthopaedic Medicine, edited by Ludwig O., Pierre B. and Herman j. 7th edition. Churchill Livingstone Publishers London.; 16:716-41.
- 13. Williams A., Newell RLM., Davies S., et al.(2005):

 The pelvic girdle and lower limb. In: Standring S, editor. 39 edition Grays Anatomy Churchill
 Livingstone Publishers, Elsevier,
 Philadelphia.; 110: 1399-1419.

- **14. Drake R., Vogl AW. & Mitchell A. (2013):** Gray's anatomy for students. Third edition. El Sevier, pp. 535-672.
- **15. Carol LA.(2000):** Evaluation of the Marrow Space in the Adult Hip. RadioGraphics; 20:S27–S42
- 16. Williams PL., Bannister LH., Berry MM., et al. (2000): Skeletal System In Gray's Anatomy, edited by Jomaah N. and Laredo JD. 38th ed. New York: Churchill Livingstone.; 55. 425-736.
- 17. David W., William JM. & James MG.(2004): The Hip. In Diagnostic Imaging Orthopaedics,. 1st edition, A Medical Reference Publishing Company, Canada.; Section 4, pp. 98.
- 18. Adam G. & Michael W.(2002): The lower limb pelvic girdle and proximal femur. In Orthopedic Radiology, edited by AdamG. and Michael W., 3rd edition Lippincott Williams & Wilkins Publishers, Philadelphia.; Section 9, 197-227.
- 19. Carol J., Shella F., Jonathan A. et al.(2003): The hip. In Essential Musculoskeletal Imaging, edited by Lee B., Mark DS., John H. et al. 1st edition. Churchill Livingstone Publishers, Philadelphia.; Section 2, 527-98.
- 20. Keith LM. & Arthur FD.(2005): Anatomy of the lower limb. In Clinically Oriented Anatomy, edited by Keith LM. and Arthur FD. 5th edition. Lippincott

- Williams & Wilkins Publishers, New York.; Section 3, 573-605.
- 21. Zlotorowicz M. & Czubak J. (2014): Vascular

 Anatomy and Blood Supply to the Femoral Head.
 in Osteonecrosis,pp.19-25.
- **22. Chung SMK. (1976):** The arterial supply of the developing proximal end of the human femur. J Bone Joint Surg Am 1976; 58(7):961-70.
- **23. Jones JP. (2000):** Etiology and pathogenesis of osteonecrosis. Sem Arthroplasty.; 2: 160-180.
- **24. Graaff VD. (2001):** Human Anatomy, Sixth Edition pp:166-219.Wilson J&, Furukawa M.: Evaluation of the patient with hip pain. Am Fam Physician. 2014 Jan 1;89(1):27-34.
- **25. William BM. & Timothy GS. (2008):** Imaging of the hip and pelvis. In first edition. Problem Solving in Musculoskeletal Imaging. Mosby/Elsevier, Chapter 11, pp. 509-564.
- **26. Lim SJ. & Park YS. (2015):** Plain Radiography of the Hip: A Review of Radiographic Techniques and Image Features. Hip & Pelvis, 27(3), 125–134.
- 27. Nestorova R., Vlad V., Petranova T., et al.(2012):

 Ultrasonography of the hip. In Medical

 Ultrasonography, 14 (3): 217-24.

- **28.** Verbeeten KM., Hermann KL., Hasselqvist M., et al (2005): The advantages of MRI in the detection of occult hip fractures. Eur Radiol 15(1):165-169.
- 29. David WS., Thomas GS., & Miriam AB. (2007): The Hip. In David W. Stoller. Third edition. Magnetic resonance imaging in orthopedics & sports medicine. Lippincott Williams & Wilkins. Pp: 42-304.
- 30. Aldridge JM. 3rd &Urbaniak JR.(2004): Avascular necrosis of the femoral head: Etiology, pathophysiology, classification, and current treatment guidelines. Am. J. Orthop. (Belle Mead NJ);33(7):327–32.
- 31. Vande Berg BC., Malghem J., Lecouvet FE., et al. (2001): Magnetic resonance imaging and differential diagnosis of epiphyseal osteonecrosis. Semin Musculoskelet Radiol.;5:57-67.
- **32. Alshahrani H. &Wood GCA.(2016):** Treatment of Avascular Necrosis of the Femoral Head with Bone Graft and Interference Screw. MOJ Orthop Rheumatol; 4(6): 00159
- **33. Takatori Y. & Ninomiya S.(2003):** Avascular necrosis of the femoral head. Natural history and magnetic resonance imaging. J Bone Joint Surg. Br.; 75(2): 217-21.

- 34. Kawashima M., Torisu T., Hayashi K., et al.(2002):

 Pathological review of osteonecrosis in divers.

 Clin Orthop.; (130): 107-17.
- 35. Manenti G., Altobelli S., Pugliese L., et al.(2015):

 The role of imaging in diagnosis and management of femoral head avascular necrosis. Clin Cases

 Miner Bone Metab.12 (Suppl 1):31-8.
- 36. Sugano N., Takaoka K., Ohzono K., et al.(1994):

 Prognostication of nontraumatic avascular
 necrosis of the femoral head. Significance of
 location and size of the necrotic lesion. Clin
 Orthop.(303):155-64.
- **37. Malizos KN., et al. (2007):** Osteonecrosis of the femoral head: Etiology, imaging and treatment. European Journal of Radiology (63) 16–28.
- **38. Pavelka K.(2000):** Osteonecrosis. *Baillieres Best Pract Res Clin Rheumatol.* 14(2):399-414. [Medline].
- **39.** James R. & Michael J.(2012): Femoral head fractures. Musculoskelet Med; 5:199–205
- **40. Thomas P. & Hans LB.(2008):** Imaging of the Musculoskeletal System; Chapter 22 page 290.
- **41. David P. (2010)**: Slipped Capital Femoral Epiphysis: Diagnosis and Management Am Fam Physician. 1;82(3):258-62.

- 42. Donald R. & Thomas GG. (2005): Traumatic Diseases; Physical Injury: Concepts and Terminology. In Donald Resnick, Mark J. Kransdorf. 3rd edition. Bone and joint imaging. Elsevier. pp. 789-830.
- 43. Raymond TM. & Stuart LW.(2006): Legg-Calve-Perthes Syndrome. In 6th Edition Lovell & Winter's Pediatric Orthopaedics; 1038-85.
- **44. Tripathy S. & Sen R. (2010):** Legg-calve-perthes Disease Current Concepts. Web. med. Central Orthopaedics;1(11).
- **45. Blumhagen J.(1994):** Legg-Calve-Perthes disease. In: Hilton SW, Edwards DK, eds. Practical Pediatric Radiology.2nd ed. Philadelphia:. WB Saunders Co.:485-96.
- **46. De Sadeleer C., Audiens H., de Boeck H., et al. (1994):** Legg-Perthes disease: partial or total loss of femoral head? Contribution of SPECT images. *Clin Nucl Med.* 19(9):830-1.
- **47. Steinberg ME. (1997):** Avascular necrosis: diagnosis, staging, and management. J Musculoskel Med. 14(11):13-25.
- **48. Chan KL. & Mok CC. (2012):** Glucocorticoid-Induced Avascular Bone Necrosis: Diagnosis and management. The Open Orthopaedics Journal, 6, 449-457.

- **49. Lavernia CJ.**, **Sierra RJ.** & **Grieco FR.(1999):**Osteonecrosis of the femoral head. J Am Acad
 Orthop Surg. 7(4):250-61.
- 50. Boechat MI., Winters WD., Hogg RJ, et al. (2001):

 Avascular necrosis of the femoral head in children with chronic renal disease. Radiology. 218(2):411-3.
- 51. Michalecki L., Gabrys D., Kulik R. et al. (2011):
 Radiotherapy induced hip joint avascular necrosis
 —two cases report. Rep Pract Oncol
 Radiother. ;16:198-201
- **52. Okane JW.(2000):** Anterior hip pain. American family physician, ; 14(5): 81-99.
- **53. Berendt T. &Byren I.(2004):** Bone and joint infection. Clin. Med.; Nov.-Dec., 4(6): 510-18.
- 54. Satomi N. &Motoki S.(2011): Seasonal Trends in the Incidence of Hip Osteoarthritis in Japanese Patients. The Open Orthopaedics Journal. 5, 134-37
- **55. Landewe R. & Heijde D.(2005):** Radiographic progression in rheumatoid arthritis. The Netherlands. Clin. Exp. Rheumatol.; Sep-Oct., 23(39): 63-68.
- **56. Zizic TM., Hungerford DS. & Stevens MB.(2002):** Ischemic bone necrosis in systemic lupus

- erythematosus. II. The early diagnosis of ischemic necrosis of bone. Medicine;59(2):134-42.
- **57. Andrea SD. & Paul B. (2009):** Imaging Investigation of Arthritis in Children. In Barbara N. Weissman. Imaging of Arthritis and Metabolic Bone Disease. Elsevier. pp 428-56.
- **58. Karagkevrekis CB. & Ainscow DA.(1998):** Transient osteoporosis of the hip associated with osteogenesis imperfecta. J Bone Joint Surg Br.;80(1):54-5.
- 59. Domachevsky L.& Keynan Y. (2004): Transient osteoporosis associated with hyperhomocystinemia: a possible role for hyperbaric oxygen therapy. Undersea Hyperb. Med.;31(3):275-9.
- **60. Stephen KS. & David LS.(2006):** Developmental Dysplasia of the Hip. Am Fam Physician ;74:1310-16
- 61. Boaz KK. & Richard BG. (2009): ACR
 Appropriateness Criteria on Developmental
 Dysplasia of the Hip-Child. J Am Coll
 Radiol.;6(8):551-57.
- **62. John DR. (2006):** Hip joint In: Edelman RR. Et al. Clinical magnetic resonance imaging 3rd ed Philadelphia, Pa: Saunders Elsevier. pp 3366-99.

- 63. Clyde AH., et al.(2009): Basic principles of musculoskeletal MRI. In: second edition. Musculoskeletal MRI. WB Saunders company, Philadelphia, pp. 1 -19.
- **64. Torsten BM. & Emil R.(2003) :** MRI Parameters and Positioning. Thieme Stuttgart New York.; 106-09.
- 65. Eva L., Pilar F., et al.(2008): Pelvis-Hip: Technical Aspects, Normal Anatomy, Common Variants, and Basic Biomechanics. In Thomas Lee Pope, Hans L. Bloem. et al. First edition. Imaging of the Musculoskeletal System. Saunders. pp 402-33.
- **66. Clyde AH., et al.(2009):** Hips and pelvis in: Musculoskeletal MRI 2nd ed. WB Saunders company, Philadelphia, pp. 324 -51.
- 67. Sarah DB. (2014): Overview of Diagnostic Imaging of Hip Joint. In Young-Jo Kim. Tallal Charles

 Mamisch. First edition. Hip Magnetic Resonance Imaging. Springer. pp 129-48.
- **68. Josef K., Gerhard L., et al. (2006):** MR and MR Arthrography. In A. M. Davies, K. Johnson et al. first edition. Imaging of the Hip & Bony Pelvis Techniques and Applications. Springer. pp 31-48.
- 69. Theysohn JM., Kraff O., Theysohn N., et al.(2014):
 Hip imaging of avascular necrosis at 7 Tesla
 compared with 3 Tesla. *Skeletal Radiol*. 43
 (5):623-32.

- 70. Zilkens C., Miese F., Jager M., et al. (2011):

 Magnetic resonance imaging of hip joint cartilage and labrum. *Orthopedic Reviews*, 3(2), e9.
- 71. Tripathi P., Singh S. & Khantal N.(2016): Hip
 Pathology Findings on Magnetic Resonance
 Imaging: A Study from Tertiary Care Institute. Int
 J Sci Stud. ;4(3):35-38
- **72. Apostolos HK. & Eleni ED. (2011):** The role of MR imaging in avascular necrosis of the femoral head. Semin Musculoskelet Radiol. 2011 Jul;15(3):281-300.
- 73. Kaushik AP., Das A. & Cui Q.(2012): Osteonecrosis of the femoral head: An update in year 2012.

 World J Orthop 2012; 3(5): 49-57.
- 74. Malizos KN., Karantanas AH., Varitimidis SE., et al. (2007): Osteonecrosis of the femoral head: Etiology, imaging and treatment. European Journal of Radiology (63) 16–28.
- 75. Mont MA., Marulanda GA., Jones LC., et al. (2006):

 Systematic analysis of classification systems for osteonecrosis of the femoral head. J Bone Joint Surg Am 88 Suppl 3:16-26.
- **76. Simone W. & Klaus W.(2014):** Measurements and Classifications in Musculoskeletal Radiology.; 39-42

- 77. Jonathan RD. & Ramiro JH.(2009): MRI of Legg-Calvé-Perthes Disease. AJR: 193:1394-1407.
- **78. Viana SL. &Machado B. (2013):** Legg-Calvé-Perthes Disease (Chapter 7). in: First edition. Joint imaging in childhood and adolescence. Springer, Heidelberg; 195–236.
- 79. Weber MA., Merle C., Rehnitz C., etal.(2016):

 Modern Radiological Imaging of Osteoarthritis of
 The Hip Joint With Consideration of Predisposing
 Conditions. Fortschr Röntgenstr; 188: 635–51.
- 80. Barbara NW. (2009): Osteoarthritis. In Barbara N. Weissman. Imaging of Arthritis and Metabolic Bone Disease. Elsevier. pp 107-33.
- **81. Herwig I. (2006):** Arthritis 1: Hip. In A. M. Davies, K. Johnson. et al. First edition. Imaging of the Hip & Bony Pelvis; Techniques and Applications. Springer. pp. 283-98.
- **82. Nathalie B., et al. (2007):** Early Rheumatoid Arthritis: A Review of MRI and Sonographic Findings. AJR; 189:1502–1509.
- **83. José AN. (2010):** MR Imaging of Early Rheumatoid Arthritis. RadioGraphics; 30:143–165.
- **84. Michael AB. (2009):** MRI In Rheumatoid Arthritis. In Michael A. Bruno, timothy J. mosher. et al. First edition. Arthritis in Color: advanced imaging of arthritis. Saunders. Philadelphia, PP 123-36.

- **85.** Manaster BJ., May DA. & Disler DG. (2007):
 Osteoporosis. In .-3rd edition. Musculoskeletal imaging. Mosby, Inc. pp 391-96.
- 86. Vande Berg BC., Lecouvet FE., Maldague B., et al. (2006): Osteonecrosis and Transient Osteoporosis of the Femoral Head, In A. M. Davies, K. Johnson. et al. First edition. Imaging of the Hip & Bony Pelvis; Techniques and Applications. Springer. pp. 195-216.
- 87. Curtis WH., William FC., et al. (1993): MR Imaging of Bone marrow edema pattern transient osteoporosis-transient bone marrow edema syndrome or osteonecrosis. RadioGraphics . 13:1001-1011.
- **88.** Luis SB., Zehava SR., et al.(2013): Imaging
 Evaluation of Developmental Hip Dysplasia in the
 Young Adult AJR; 200:1077-88.
- **89. Starr V. & Bo YH.(2014):** Imaging update on developmental dysplasia of the hip with the role of MRI. AJR; 203:1324–35.
- **90. Mayumi Oka. & Johnny UVM.(2004):** Prevalence and Patterns of Occult Hip Fractures and Mimics Revealed by MRI. AJR ;182:283–288.
- **91. Takuaki Y.(2012):** Subchondral Insufficiency Fractures of the Femoral Head Clinics in Orthopedic Surgery ;4:173-80.

- 92. José MM., Ana MH., et al.(2008): Acute Osseous
 Injury to the Hip and Proximal Femur. In Thomas
 Pope, Hans L. Bloem et al. first edition. Imaging
 of the Musculoskeletal System. Saunders.
 Philadelphia, pp 470-98.
- 93. Jeffrey JN. & Young JK.(2014): Developmental Hip Disorders. In Young-Jo Kim. Tallal Charles Mamisch. First edition. Hip Magnetic Resonance Imaging. Springer. pp 95-112.
- 94. Jaimes C., Chauvin NA., Delgado J., et al.(2014):

 MR imaging of normal epiphyseal development
 and common epiphyseal disorders.

 Radiographics;34(2):449-71.
- 95. Nori M., Marupaka SK., Alluri S., et al. (2015): MRI Evaluation of Post Core Decompression Changes in Avascular Necrosis of Hip. Journal of Clinical and Diagnostic Research: 9(12), p. 4-8.
- 96. Horia B., Valentin C. & Mihaela D.(2009):

 Avascular necrosis of the femoral head. journal of clinical medicine, volume 4 No.1: 26-34.
- 97. Hamilton TW., Goodman SM., & Figgie M. (2009): SAS Weekly Rounds: Avascular Necrosis. HSS Journal, 5(2), 99-113.
- 98. Manenti G., Altobelli S., Pugliese L., et al.(2015):

 The role of imaging in diagnosis and management of femoral head avascular necrosis. Clinical Cases

in Mineral and Bone Metabolism, 12(Suppl 1), 31–38.

- 99. Glickstein MF., Burk DL., Schiebler ML., et al. (1988): Avascular necrosis versus other diseases of the hip: sensitivity of MR imaging. Radiology. 169 (1): 213-5.
- 100. Steinberg ME., Hayken GD, Steinberg DR.(1995):
 A quantitative system for staging avascular
 necrosis. J Bone Joint Surg Br. 77:34-41.
- 101. Tripathy SK., Goyal T., & Sen RK. (2015):

 Management of femoral head osteonecrosis:

 Current concepts. Indian Journal of

 Orthopaedics, 49(1), 28-45.
- 102. Doré D., de Hoog J., Giles G., et al (2012): A longitudinal study of the association between dietary factors, serum lipids, and bone marrow lesions of the knee. Arthritis Research & Therapy, 14(1), R13.
- 103. Teichtahl AJ., Wang Y., Smith S., et al.
 (2014):Structural changes of hip osteoarthritis
 using magnetic resonance imaging. Arthritis Res
 Ther. 16:466.
- 104. Xu L., Hayashi D., Roemer FW., Felson DT., et al.(2012): Magnetic Resonance Imaging ofSubchondral Bone Marrow Lesions in Association

with Osteoarthritis. Seminars in Arthritis and Rheumatism, 42(2), 105–118.

- 105. Boutry N., Khalil C., Jaspart M., et al.(2007):
 Imaging
 of the hip in patients with rheumatic disorders.
 Eur J Radiol. 63(1):49-58.
- 106. Jacobson JA., Girish G., Jiang Y., et al.(2008):

 Radiographic evaluation of arthritis: degenerative joint disease and variations.

 Radiology. 248(3):737-47
- **107. McQueen FM. (2000):** Magnetic resonance imaging in early inflammatory arthritis: what is its role?. Rheumatology (Oxford). 39(7):700-6.
- **108. Braun HJ. & Gold GE. (2012):** Diagnosis of Osteoarthritis: Imaging. Bone, 51(2), 278–288.
- 109. Van WK., Pritchard P., & Taylor JA. (2013):

 Transient osteoporosis of the hip: A case
 report. The Journal of the Canadian Chiropractic
 Association, 57(2), 116–122.
- **110. McWalter P., & Hassan A. (2009):** Transient osteoporosis of the hip. Annals of Saudi Medicine, 29(2), 146–148.
- 111. Mirza R., Ishaq S. & Amjad H.(2012): Transient osteoporosis of the hip. J Pak Med Assoc. Feb;62(2):196-8.

- **112. Balakrishnan A., Schemitsch EH., Pearce D., et al. (2003):** Distinguishing transient osteoporosis of the hip from avascular necrosis. Canadian Journal of Surgery, 46(3), 187-192.
- 113. Karchevsky M., Schweitzer ME., Morrison WB., et al.(2004): MRI findings of septic arthritis and associated osteomyelitis in adults. AJR Am J Roentgenol. 182(1):119-22.
- **114. Learch TJ. & Farooki S. (2006):** Magnetic resonance imaging of septic arthritis. Clin Imaging; 24:236 -242.
- 115. Sherwani P., Vire A., Anand R., et al.(2016):

 Utility of MR Imaging in Developmental Dysplasia of Hip. Austin J Radiol. 3(3): 1052.
- 116. Evan JZ., Edward YL., Ricardo R., et al.(2013):
 Hip Disorders in Children AJR. 201:W776-W796.
- 117. Jaramillo D., Galen TA., Winalski CS., et al.
 (1999): Legg-Calvé-Perthes disease: MR imaging evaluation during manual positioning of the hip—comparison with conventional arthrography.
 Radiology. 212:519-25.
- **118. Dwek JR.(2009):** The hip: MR imaging of uniquely pediatric disorders. Magn Reson Imaging Clin N Am. 17(3):509-20.
- 119. Lahdes VT., Lamminen A., Merikanto J., et al.(1997): The value of MRI in early Perthes'

disease: An MRI study with a 2-year follow-up. Pediatr Radiol. 27(6):517-22.

120. Staatz C., Honnef D., Kochs A., et al.(2007):

Evaluation of femoral head vascularization in slipped capital femoral epiphysis before and after cannulated screw fixation with use of contrast-enhanced MRI initial results. Eur Radiol. 17(1):163-8.

121. Green DW., Reynolds RAK., Khan SN., et al.

(2005): The Delay in Diagnosis of Slipped Capital Femoral Epiphysis: A Review of 102 Patients. HSS Journal, 1(1), 103–106.

122. Propeck T. & Ridpath CA.(2001): Slipped capital femoral epiphysis diagnosed by magnetic resonance imaging. Emergency Radiology. 8: 344-46.

123. Dillon JE., Connolly SA. & Connolly LP. (2009):

MR Imaging of Congenital/Developmental and Acquired Disorders of the Pediatric Hip and Pelvis. Magnetic Resonance Imaging Clinics of North America, 13: 55-80.

124. Anand A., RaviRaj A. & Kodikal G. (2010):

Subchondral stress fracture of femoral head in a healthy adult. Indian Journal of Orthopaedics, 44(4), 458-60.

125. Kimura T., Goto T., Hamada D., et al.(2016):

Subchondral Insufficiency Fracture of the Femoral Head Caused by Excessive Lateralization of the Acetabular Rim. Case Reports in Orthopedics, vol. 2016, Article ID 4371679, 5 pages.

- 126. Hiren P. & Atul FK. (2016): Subchondral
 Insufficiency Fracture of the Femoral Head
 treated with Core Decompression and Bone Void
 Filler Support. Arch Bone Jt Surg. 4(3): 264-68.
- 127. Ikemura S., Yamamoto T., Motomura G., et al. (2010): MRI evaluation of collapsed femoral heads in patients 60 years old or older: differentiation of subchondral insufficiency fracture from osteonecrosis of the femoral head. AJR Am J Roentgenol. 195(1):W63-W68.

